

Computer-aided proofs in dynamics I

Interval arithmetic and maps

Maciej Capiński

AGH University of Kraków

Plan of the lecture

- Computer-aided proofs - “philosophy”
- Brouwer theorem
- Interval Newton/Krawczyk method
- Wrapping effect
- Local coordinates


Literature

- W. Tucker, *Validated Numerics: A Short Introduction to Rigorous Computations*, Princeton University Press 2011
- J-B van den Berg *Introduction to rigorous numerics in dynamics: general functional analytic setup and an example that forces chaos* Proc. Sympos. Appl. Math. 2018
- T. Kapela, M. Mrozek, D. Wilczak, P. Zgliczyński, CAPD::DynSys: a flexible C++ toolbox for rigorous numerical analysis of dynamical systems, Commun. Nonlinear Sci. Numer. Simul. 2021

Computer-aided proofs - philosophy

Bolzano theorem

$$f : \mathbb{R} \rightarrow \mathbb{R} \quad f(x) \stackrel{?}{=} 0$$



$$f(a) < 0 \quad f(b) > 0$$


- There exists an x^* in (a, b) such that

$$f(x^*) = 0$$

Computer-aided proofs - philosophy

Bolzano theorem - no need to be too accurate

$$f : \mathbb{R} \rightarrow \mathbb{R} \qquad f(x) \stackrel{?}{=} 0$$



$$f(a) < 0 \qquad f(b) > 0$$

- There exists an x^* in (a, b) such that


$$f(x^*) = 0$$

Computer-aided proofs - philosophy

Bolzano theorem - no need to be too accurate

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) \stackrel{?}{=} 0$$



$$f(a) < 0 \quad f(b) > 0$$

CAP- the ingredients:

- Mathematical theorem
- Good understanding of the numerical properties of the problem
- Computer-aided validation of assumptions

- There exists an x^* in (a, b) such that

$$f(x^*) = 0$$

Kinds of tools that we shall use

Interval arithmetic

computations on intervals:

$$[1, 2] + [3, 4] = [4, 6]$$

$$[1, 2] - [3, 4] = [-3, -1]$$

$$[1, 2] * [3, 4] = [3, 8]$$


$$[1, 2] / [3, 4] = \left[\frac{1}{4}, \frac{2}{3} \right]$$

$$[1, 2]^{[3,4]} = [1^3, 2^4]$$

...

extends to higher dimensions

$$[1, 2] - [1, 2] = [-1, 1]$$



$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

What can be computed:

- $[f(U)]$, $[Df(U)]$
- higher order derivatives
- linear algebra; eg. $[A^{-1}]$

INTLAB^a

RadiiPolynomial^b

CAPD^c

^awww.tuhh.de/ti3/rump/intlab

^bgithub.com/OlivierHnt/RadiiPolynomial.jl

^ccapd.ii.uj.edu.pl

Kinds of tools that we shall use

Interval arithmetic

computations on intervals:

$$[1, 2] + [3, 4] = [4, 6]$$

$$[1, 2] - [3, 4] = [-3, -1]$$

$$[1, 2] * [3, 4] = [3, 8]$$


$$[1, 2] / [3, 4] = \left[\frac{1}{4}, \frac{2}{3} \right]$$

$$[1, 2]^{[3,4]} = [1^3, 2^4]$$

...

extends to higher dimensions

$$[1, 2] - [1, 2] = [-1, 1]$$



$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

What can be computed:

- $[f(U)]$, $[Df(U)]$
- higher order derivatives
- linear algebra; eg. $[A^{-1}]$

INTLAB^a

RadiiPolynomial^b

CAPD^c

^awww.tuhh.de/ti3/rump/intlab

^bgithub.com/OlivierHnt/RadiiPolynomial.jl

^ccapd.ii.uj.edu.pl

Kinds of tools that we shall use

Interval arithmetic

computations on intervals:

$$[1, 2] + [3, 4] = [4, 6]$$

$$[1, 2] - [3, 4] = [-3, -1]$$

$$[1, 2] * [3, 4] = [3, 8]$$


$$[1, 2] / [3, 4] = \left[\frac{1}{4}, \frac{2}{3} \right]$$

$$[1, 2]^{[3,4]} = [1^3, 2^4]$$

...

extends to higher dimensions

$$[1, 2] - [1, 2] = [-1, 1]$$



$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

What can be computed:

- $[f(U)]$, $[Df(U)]$
- higher order derivatives
- linear algebra; eg. $[A^{-1}]$

INTLAB^a

RadiiPolynomial^b

CAPD^c

^awww.tuhh.de/ti3/rump/intlab

^bgithub.com/OlivierHnt/RadiiPolynomial.jl

^ccapd.ii.uj.edu.pl


Examples of tools that we can use

Let $B \subset \mathbb{R}^n$ be homeomorphic to a closed ball

Theorem (Brouwer theorem)

If $F : B \rightarrow B$ is continuous, then there exists a $q \in B$

$$F(q) = q$$



Examples of tools that we can use


Theorem (Banach fixed pt thm)

Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$. If for some $\lambda < 1$

$$\|F(x_1) - F(x_2)\| \leq \lambda \|x_1 - x_2\|$$

then there exists a q such that

$$F(q) = q$$




- The use of contraction can be exploited in various contexts

Interval Newton method

$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad C^1$$

$$B = \prod_{i=1}^n [a_i, b_i]$$

$$x_0 \in B$$



Theorem (interval Newton)

If

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B$$

Then $\exists! x^* \in B$ such that

$$F(x^*) = 0$$

(intuition)
Newton-Raphson:

$$F : \mathbb{R} \rightarrow \mathbb{R}$$

$$x_{n+1} = x_n - \frac{F(x_n)}{F'(x_n)}$$


Proof. chalk.

Interval Newton method

$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad C^1$$

$$B = \prod_{i=1}^n [a_i, b_i]$$

$$x_0 \in B$$



Theorem (interval Newton)

If

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B$$

Then $\exists! x^* \in B$ such that

$$F(x^*) = 0$$

(intuition)

Newton-Raphson:

$$F : \mathbb{R} \rightarrow \mathbb{R}$$

$$x_{n+1} = x_n - \frac{F(x_n)}{F'(x_n)}$$


Proof. chalk.

Interval Newton method

$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad C^1$$

$$B = \prod_{i=1}^n [a_i, b_i]$$

$$x_0 \in B$$



Theorem (interval Newton)

If

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B$$

Then $\exists! x^* \in B$ such that

$$F(x^*) = 0$$

(intuition)

Newton-Raphson:

$$F : \mathbb{R} \rightarrow \mathbb{R}$$

$$x_{n+1} = x_n - \frac{F(x_n)}{F'(x_n)}$$


Proof. chalk.

Interval Newton method

$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad C^1$$

$$B = \prod_{i=1}^n [a_i, b_i]$$

$$x_0 \in B$$



Theorem (interval Newton)

If

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B$$

Then $\exists! x^* \in B$ such that

$$F(x^*) = 0$$

(intuition)

Newton-Raphson:

$$F : \mathbb{R} \rightarrow \mathbb{R}$$

$$x_{n+1} = x_n - \frac{F(x_n)}{F'(x_n)}$$


Proof. chalk.

Interval Newton method

$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad C^1$$

$$B = \prod_{i=1}^n [a_i, b_i]$$

$$x_0 \in B$$



Theorem (interval Newton)

If

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B$$

Then $\exists! x^* \in B$ such that

$$F(x^*) = 0$$

(intuition)
Newton-Raphson:

$$F : \mathbb{R} \rightarrow \mathbb{R}$$

$$x_{n+1} = x_n - \frac{F(x_n)}{F'(x_n)}$$


Proof. chalk.

Interval Newton method

$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad C^1$$

$$B = \prod_{i=1}^n [a_i, b_i]$$

$$x_0 \in B$$



Theorem (interval Newton)

If

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B$$

Then $\exists! x^* \in B$ such that

$$F(x^*) = 0$$

(intuition)
Newton-Raphson:

$$F : \mathbb{R} \rightarrow \mathbb{R}$$

$$x_{n+1} = x_n - \frac{F(x_n)}{F'(x_n)}$$

Proof. chalk.

Interval Newton method

Example - Hénon map

Theorem (interval Newton)

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B.$$

Then $\exists! x^* \in B, \quad F(x^*) = 0.$

$$h(x, y) = (1 - ax^2 + y, bx)$$

$$a = 1.4, b = 0.3$$

Fixed point:

$$h(x, y) = (x, y)$$

$$F(x, y) = (1 - ax^2 + y - x, bx - y) = 0$$



Interval Newton method

Example - Hénon map

Theorem (interval Newton)

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B.$$

Then $\exists! x^* \in B, \quad F(x^*) = 0.$

$$h(x, y) = (1 - ax^2 + y, bx)$$

$$a = 1.4, b = 0.3$$

Fixed point:

$$h(x, y) = (x, y)$$

$$F(x, y) = (1 - ax^2 + y - x, bx - y) = 0$$



Interval Newton method

Example - Hénon map

Theorem (interval Newton)

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B.$$

Then $\exists! x^* \in B, \quad F(x^*) = 0.$

$$h(x, y) = (1 - ax^2 + y, bx)$$

$$a = 1.4, b = 0.3$$

Fixed point:

$$h(x, y) = (x, y)$$

$$F(x, y) = (1 - ax^2 + y - x, bx - y) = 0$$



Interval Newton method

Example - Hénon map

Theorem (interval Newton)

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B.$$

Then $\exists! x^* \in B, \quad F(x^*) = 0.$

$$F(x, y) = (1 - ax^2 + y - x, bx - y) = 0$$

```
IMap F = "par:a,b;  
         var:x,y;  
         fun:1-a*x^2+y-x,b*x-y;"
```

```
IVector x0(2);  
x0[0] = sqrt(609)/28-0.25;  
x0[1] = b*x0[0];
```

```
IVector B(2);  
B[0] = x0[0] + interval(-1,1)*power(10,-14);  
B[1] = x0[1] + interval(-1,1)*power(10,-14);
```

```
IVector N = x0 - gauss( F[B] , F(x0) );
```

```
if( subsetInterior(N,B) ) return 1;
```



(show Newton method program)

Interval Newton method

Example - Hénon map

Theorem (interval Newton)

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B.$$

Then $\exists! x^* \in B, \quad F(x^*) = 0.$

$$F(x, y) = (1 - ax^2 + y - x, bx - y) = 0$$

```
IMap F = "par:a,b;  
         var:x,y;  
         fun:1-a*x^2+y-x,b*x-y";
```

```
IVector x0(2);  
x0[0] = sqrt(609)/28-0.25;  
x0[1] = b*x0[0];
```

```
IVector B(2);  
B[0] = x0[0] + interval(-1,1)*power(10,-14);  
B[1] = x0[1] + interval(-1,1)*power(10,-14);
```

```
IVector N = x0 - gauss( F[B] , F(x0) );
```

```
if( subsetInterior(N,B) ) return 1;
```



(show Newton method program)

Interval Newton method

Example - Hénon map

Theorem (interval Newton)

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B.$$

Then $\exists! x^* \in B, \quad F(x^*) = 0.$

$$F(x, y) = (1 - ax^2 + y - x, bx - y) = 0$$

```
IMap F = "par:a,b;  
         var:x,y;  
         fun:1-a*x^2+y-x,b*x-y";
```

```
IVector x0(2);  
x0[0] = sqrt(609)/28-0.25;  
x0[1] = b*x0[0];
```

```
IVector B(2);  
B[0] = x0[0] + interval(-1,1)*power(10,-14);  
B[1] = x0[1] + interval(-1,1)*power(10,-14);
```

```
IVector N = x0 - gauss( F[B] , F(x0) );
```

```
if( subsetInterior(N,B) ) return 1;
```



(show Newton method program)

Interval Newton method

Example - Hénon map

Theorem (interval Newton)

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B.$$

Then $\exists! x^* \in B, \quad F(x^*) = 0.$

$$F(x, y) = (1 - ax^2 + y - x, bx - y) = 0$$

```
IMap F = "par:a,b;  
         var:x,y;  
         fun:1-a*x^2+y-x,b*x-y;"
```

```
IVector x0(2);  
x0[0] = sqrt(609)/28-0.25;  
x0[1] = b*x0[0];
```

```
IVector B(2);  
B[0] = x0[0] + interval(-1,1)*power(10,-14);  
B[1] = x0[1] + interval(-1,1)*power(10,-14);
```

```
IVector N = x0 - gauss( F[B] , F(x0) );
```

```
if( subsetInterior(N,B) ) return 1;
```

(show Newton method program)



Interval Newton method

Example - Hénon map

Theorem (interval Newton)

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B.$$

Then $\exists! x^* \in B, \quad F(x^*) = 0.$

$$F(x, y) = (1 - ax^2 + y - x, bx - y) = 0$$

```
IMap F = "par:a,b;  
         var:x,y;  
         fun:1-a*x^2+y-x,b*x-y;";  
  
IVector x0(2);  
x0[0] = sqrt(609)/28-0.25;  
x0[1] = b*x0[0];  
  
IVector B(2);  
B[0] = x0[0] + interval(-1,1)*power(10,-14);  
B[1] = x0[1] + interval(-1,1)*power(10,-14);  
  
IVector N = x0 - gauss( F[B] , F(x0) );  
  
if( subsetInterior(N,B) ) return 1;
```



(show Newton method program)

Interval Newton method

Example - Hénon map

Theorem (interval Newton)

$$x_0 - [DF(B)]^{-1}F(x_0) \subset B.$$

Then $\exists! x^* \in B, \quad F(x^*) = 0.$

$$F(x, y) = (1 - ax^2 + y - x, bx - y) = 0$$


```
IMap F = "par:a,b;  
         var:x,y;  
         fun:1-a*x^2+y-x,b*x-y;";  
  
IVector x0(2);  
x0[0] = sqrt(609)/28-0.25;  
x0[1] = b*x0[0];  
  
IVector B(2);  
B[0] = x0[0] + interval(-1,1)*power(10,-14);  
B[1] = x0[1] + interval(-1,1)*power(10,-14);  
  
IVector N = x0 - gauss( F[B] , F(x0) );  
  
if( subsetInterior(N,B) ) return 1;
```



(show Newton method program)

Krawczyk Theorem

To get better results one gets creative



Theorem

Let

$$K = x_0 - CF(x_0) + (\text{Id} - C[DF(B)])(B - x_0)$$

If $K \subset \text{int}(B)$ then $\exists! x^* \in B$ such that

$$F(x^*) = 0$$

$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$$B = \Pi_{i=1}^n [a_i, b_i]$$

$$x_0 \in B$$

$$C \in \mathbb{R}^{n \times n}$$


- $F(x_0) \approx 0$
- $C \approx DF(x_0)^{-1}$

(show Krawczyk method program)

Krawczyk vs Newton

- $x_0 \in B$
- $F(x_0) \approx 0$

- $C \approx DF(x_0)^{-1}$



Newton method

$$N = x_0 - [DF(B)]^{-1}F(x_0)$$

$$N \subset B?$$

- solving a linear problem in interval arithmetic

Krawczyk method

$$K = x_0 - CF(x_0) + (\text{Id} - C[DF(B)])(B - x_0)$$


$$K \subset \text{int}(B)?$$

- guessing/computing C non-rigorously

Krawczyk vs Newton

- $x_0 \in B$
- $F(x_0) \approx 0$

- $C \approx DF(x_0)^{-1}$



Newton method

$$N = x_0 - [DF(B)]^{-1}F(x_0)$$

$$N \subset B?$$

- solving a linear problem in **interval arithmetic**

Krawczyk method

$$K = x_0 - CF(x_0) \\ + (\text{Id} - C[DF(B)])(B - x_0)$$

$$K \subset \text{int}(B)?$$

- guessing/computing C **non-rigorously**


Fork in the road

Theorem (Banach fixed pt thm)

Let $F : X \rightarrow X$. If for some $\lambda < 1$

$$\|F(x_1) - F(x_2)\| \leq \lambda \|x_1 - x_2\|$$

then there exists a q such that
 $F(q) = q$.



Analytic approach

- Solving ODEs
- Boundary value problems
- Invariant manifolds
- DDEs, PDEs

geometric & topological approach

- Chaotic dynamics for ODEs
- Invariant manifolds
- **Blenders**
- DDEs, PDEs

J-B van den Berg "Introduction to rigorous numerics in dynamics: general functional analytic setup" <https://www.math.vu.nl/~janbouwe/pub/introrignumdyn.pdf>

Periodic orbits

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$


$$f^k(x) = x \text{ when}$$

$$F(x) = 0$$

$$\text{for } F(x) = f^k(x) - x$$

$$f(x) = Rx + g(x)$$


R is rotation



Problem with Newton method

$$N = x_0 - [DF(B)]^{-1}F(x_0)$$

$$[Df^k(B)] = [Df(f^{k-1}(B))] \cdot [Df^{k-1}(B)]$$



$$B_0 = B \quad D_0 = \text{Id}$$

For $i = 1, \dots, k$

$$B_i = [f(B_{i-1})] \quad D_i = [Df(B_{i-1})] \cdot D_{i-1}$$

Then

$$[Df^k(B)] = D_k$$

Periodic orbits

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$


$$f^k(x) = x \text{ when}$$

$$F(x) = 0$$

$$\text{for } F(x) = f^k(x) - x$$

$$f(x) = Rx + g(x)$$


R is rotation



Problem with Newton method

$$N = x_0 - [DF(B)]^{-1}F(x_0)$$

$$[Df^k(B)] = [Df(f^{k-1}(B))] \cdot [Df^{k-1}(B)]$$



$$B_0 = B \quad D_0 = \text{Id}$$

For $i = 1, \dots, k$

$$B_i = [f(B_{i-1})] \quad D_i = [Df(B_{i-1})] \cdot D_{i-1}$$

Then

$$[Df^k(B)] = D_k$$

Periodic orbits

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$


$$f^k(x) = x \text{ when}$$

$$F(x) = 0$$

$$\text{for } F(x) = f^k(x) - x$$

$$f(x) = Rx + g(x)$$


R is rotation



Problem with Newton method

$$N = x_0 - [DF(B)]^{-1}F(x_0)$$

$$[Df^k(B)] = [Df(f^{k-1}(B))] \cdot [Df^{k-1}(B)]$$



$$B_0 = B \quad D_0 = \text{Id}$$

For $i = 1, \dots, k$

$$B_i = [f(B_{i-1})] \quad D_i = [Df(B_{i-1})] \cdot D_{i-1}$$

Then

$$[Df^k(B)] = D_k$$

Periodic orbits

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$


$$f^k(x) = x \text{ when}$$

$$F(x) = 0$$

$$\text{for } F(x) = f^k(x) - x$$

$$f(x) = Rx + g(x)$$


R is rotation



Problem with Newton method

$$N = x_0 - [DF(B)]^{-1}F(x_0)$$

$$[Df^k(B)] = [Df(f^{k-1}(B))] \cdot [Df^{k-1}(B)]$$



$$B_0 = B \quad D_0 = \text{Id}$$

For $i = 1, \dots, k$

$$B_i = [f(B_{i-1})] \quad D_i = [Df(B_{i-1})] \cdot D_{i-1}$$

Then

$$[Df^k(B)] = D_k$$

Periodic orbits

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$


$$f^k(x) = x \text{ when}$$

$$F(x) = 0$$

$$\text{for } F(x) = f^k(x) - x$$

$$f(x) = Rx + g(x)$$


R is rotation



Problem with Newton method

$$N = x_0 - [Df(B)]^{-1}F(x_0)$$

$$[Df^k(B)] = [Df(f^{k-1}(B))] \cdot [Df^{k-1}(B)]$$



$$B_0 = B \quad D_0 = \text{Id}$$

For $i = 1, \dots, k$


$$B_i = [f(B_{i-1})] \quad D_i = [Df(B_{i-1})] \cdot D_{i-1}$$

Then

$$[Df^k(B)] = D_k$$

Wrapping effect

Remedy 1 - parallel shooting



Problem with Newton method

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$$F(x) = f^k(x) - x$$

$$N = x_0 - [DF(B)]^{-1}F(x_0)$$

$$[Df^k(B)] = [Df(f^{k-1}(B))] \cdot [Df^{k-1}(B)]$$

Remedy:

$f^k(x_1) = x_1$ when $F(x) = 0$ for


$$F(x_1, \dots, x_k) = \begin{pmatrix} f(x_1) - x_2 \\ f(x_2) - x_3 \\ \vdots \\ f(x_k) - x_1 \end{pmatrix}$$

There is no free lunch:

- $F : \mathbb{R}^{nk} \rightarrow \mathbb{R}^{nk}$ instead of \mathbb{R}^n
- The matrix DF is cumbersome to compute
- Use Krawczyk method!

Wrapping effect

Remedy 2 - Take care of coordinates



$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Let us represent a set as:

$$X = x + AY$$

where $x, Y \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$

$$\begin{aligned} F(x + y) &= F(x) + \int_0^1 \frac{d}{ds} F(x + sy) ds \\ &= F(x) + \int_0^1 DF(x + sy) ds y \\ &\in F(x) + [DF(X)]y \end{aligned}$$

$$\begin{aligned} F(x + AY) &\in F(x) + ([DF(X)]A)Y \\ &= F(x) + B(B^{-1}[DF(X)]A)Y \end{aligned}$$


so $F(X) \subset \bar{X}$ for

$$\bar{X} = \bar{x} + B\bar{Y}$$

$$\bar{x} = F(x) \text{ and } \bar{Y} = (B^{-1}[DF(X)]A)Y$$

Wrapping effect

Remedy 2 - Take care of coordinates



$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Let us represent a set as:

$$X = x + AY$$

where $x, Y \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$

$$\begin{aligned} F(x + y) &= F(x) + \int_0^1 \frac{d}{ds} F(x + sy) ds \\ &= F(x) + \int_0^1 DF(x + sy) ds y \\ &\in F(x) + [DF(X)]y \end{aligned}$$

$$\begin{aligned} F(x + AY) &\in F(x) + ([DF(X)]A)Y \\ &= F(x) + B(B^{-1}[DF(X)]A)Y \end{aligned}$$


so $F(X) \subset \bar{X}$ for

$$\bar{X} = \bar{x} + B\bar{Y}$$

$$\bar{x} = F(x) \text{ and } \bar{Y} = (B^{-1}[DF(X)]A)Y$$

Wrapping effect

Remedy 2 - Take care of coordinates



$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Let us represent a set as:

$$X = x + AY$$

where $x, Y \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$

$$\begin{aligned} F(x + y) &= F(x) + \int_0^1 \frac{d}{ds} F(x + sy) ds \\ &= F(x) + \int_0^1 DF(x + sy) ds y \\ &\in F(x) + [DF(X)]y \end{aligned}$$

$$\begin{aligned} F(x + AY) &\in F(x) + ([DF(X)]A)Y \\ &= F(x) + B(B^{-1}[DF(X)]A)Y \end{aligned}$$


so $F(X) \subset \bar{X}$ for

$$\bar{X} = \bar{x} + B\bar{Y}$$

$$\bar{x} = F(x) \text{ and } \bar{Y} = (B^{-1}[DF(X)]A)Y$$

Wrapping effect

Remedy 2 - Take care of coordinates



$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Let us represent a set as:

$$X = x + AY$$

where $x, Y \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$

$$\begin{aligned} F(x + y) &= F(x) + \int_0^1 \frac{d}{ds} F(x + sy) ds \\ &= F(x) + \int_0^1 DF(x + sy) ds y \\ &\in F(x) + [DF(X)]y \end{aligned}$$

$$\begin{aligned} F(x + AY) &\in F(x) + ([DF(X)]A)Y \\ &= F(x) + B(B^{-1}[DF(X)]A)Y \end{aligned}$$


so $F(X) \subset \bar{X}$ for

$$\bar{X} = \bar{x} + B\bar{Y}$$

$$\bar{x} = F(x) \text{ and } \bar{Y} = (B^{-1}[DF(X)]A)Y$$

Wrapping effect

Remedy 2 - Take care of coordinates



$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Let us represent a set as:

$$X = x + AY$$

where $x, Y \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$

$$\begin{aligned} F(x + y) &= F(x) + \int_0^1 \frac{d}{ds} F(x + sy) ds \\ &= F(x) + \int_0^1 DF(x + sy) ds y \\ &\in F(x) + [DF(X)]y \end{aligned}$$

$$\begin{aligned} F(x + AY) &\in F(x) + ([DF(X)]A)Y \\ &= F(x) + B(B^{-1}[DF(X)]A)Y \end{aligned}$$


so $F(X) \subset \bar{X}$ for

$$\bar{X} = \bar{x} + B\bar{Y}$$

$$\bar{x} = F(x) \text{ and } \bar{Y} = (B^{-1}[DF(X)]A)Y$$

Wrapping effect

Remedy 2 - Take care of coordinates



$$F : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Let us represent a set as:

$$X = x + AY$$

where $x, Y \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$

$$\begin{aligned} F(x + y) &= F(x) + \int_0^1 \frac{d}{ds} F(x + sy) ds \\ &= F(x) + \int_0^1 DF(x + sy) ds y \\ &\in F(x) + [DF(X)]y \end{aligned}$$

$$\begin{aligned} F(x + AY) &\in F(x) + ([DF(X)]A)Y \\ &= F(x) + B(B^{-1}[DF(X)]A)Y \end{aligned}$$

so $F(X) \subset \bar{X}$ for

$$\bar{X} = \bar{x} + B\bar{Y}$$

$$\bar{x} = F(x) \text{ and } \bar{Y} = (B^{-1}[DF(X)]A)Y$$

Wrapping effect

Remedy 2 - Take care of coordinates

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Let us represent a set as:

$$X = x + AY$$

where $x, Y \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$

Example

$$f(x) = Rx + \varepsilon g(x)$$

where R is a rotation matrix

(show code)

$$f(X) \subset \bar{X} \text{ for}$$

$$\bar{X} = \bar{x} + B\bar{Y}$$

$$\bar{x} = f(x) \text{ and } \bar{Y} = (B^{-1}[Df(X)]A)Y$$

Initial set:

$$X_0 = x_0 + Y_0$$

choose some local coordinates:

$$A_k = R^k$$


$$x_k = f^k(x_0)$$

Compute inductively:

$$Y_{k+1} = (A_{k+1}^{-1} Df(X_k) A_k) Y_k$$

$$X_{k+1} = x_k + A_k Y_k$$

Then $f(X_k) \subset X_{k+1}$.



Wrapping effect

Remedy 2 - Take care of coordinates

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Let us represent a set as:

$$X = x + AY$$

where $x, Y \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$


Example

$$f(x) = Rx + \varepsilon g(x)$$

where R is a rotation matrix

(show code)

$$f(X) \subset \bar{X} \text{ for}$$



$$\bar{X} = \bar{x} + B\bar{Y}$$

$$\bar{x} = f(x) \text{ and } \bar{Y} = (B^{-1}[Df(X)]A)Y$$

Initial set:

$$X_0 = x_0 + Y_0$$

choose some local coordinates:

$$A_k = R^k$$

$$x_k = f^k(x_0)$$

Compute inductively:

$$Y_{k+1} = (A_{k+1}^{-1} Df(X_k) A_k) Y_k$$

$$X_{k+1} = x_k + A_k Y_k$$

Then $f(X_k) \subset X_{k+1}$.

Wrapping effect

Remedy 2 - Take care of coordinates

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Let us represent a set as:

$$X = x + AY$$

where $x, Y \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$


Example

$$f(x) = Rx + \varepsilon g(x)$$

where R is a rotation matrix

(show code)

$$f(X) \subset \bar{X} \text{ for}$$



$$\bar{X} = \bar{x} + B\bar{Y}$$

$$\bar{x} = f(x) \text{ and } \bar{Y} = (B^{-1}[Df(X)]A)Y$$

Initial set:

$$X_0 = x_0 + Y_0$$

choose some local coordinates:

$$A_k = R^k$$

$$x_k = f^k(x_0)$$

Compute inductively:

$$Y_{k+1} = (A_{k+1}^{-1} Df(X_k) A_k) Y_k$$

$$X_{k+1} = x_k + A_k Y_k$$

Then $f(X_k) \subset X_{k+1}$.

Wrapping effect

Remedy 2 - Take care of coordinates

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Let us represent a set as:

$$X = x + AY$$

where $x, Y \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$


Example

$$f(x) = Rx + \varepsilon g(x)$$

where R is a rotation matrix

(show code)

$$f(X) \subset \bar{X} \text{ for}$$



$$\bar{X} = \bar{x} + B\bar{Y}$$

$$\bar{x} = f(x) \text{ and } \bar{Y} = (B^{-1}[Df(X)]A)Y$$

Initial set:

$$X_0 = x_0 + Y_0$$

choose some local coordinates:

$$A_k = R^k$$

$$x_k = f^k(x_0)$$

Compute inductively:

$$Y_{k+1} = (A_{k+1}^{-1} Df(X_k) A_k) Y_k$$

$$X_{k+1} = x_k + A_k Y_k$$

Then $f(X_k) \subset X_{k+1}$.

Wrapping effect

Remedy 2 - Take care of coordinates

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Let us represent a set as:

$$X = x + AY$$

where $x, Y \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$


Example

$$f(x) = Rx + \varepsilon g(x)$$

where R is a rotation matrix

(show code)

$$f(X) \subset \bar{X} \text{ for}$$



$$\bar{X} = \bar{x} + B\bar{Y}$$

$$\bar{x} = f(x) \text{ and } \bar{Y} = (B^{-1}[Df(X)]A)Y$$

Initial set:

$$X_0 = x_0 + Y_0$$

choose some local coordinates:

$$A_k = R^k$$

$$x_k = f^k(x_0)$$

Compute inductively:


$$Y_{k+1} = (A_{k+1}^{-1} Df(X_k) A_k) Y_k$$

$$X_{k+1} = x_k + A_k Y_k$$

Then $f(X_k) \subset X_{k+1}$.

Key points

- Good numerical understanding first
- Good theorem (eg Newton vs Krawczyk)
- (Most) often being aware of coordinates helps



Tomorrow: CAP – topological shadowing