

# Computer-aided proofs in dynamics III

Blenders

Maciej Capiński

AGH University of Kraków

joint work with

Bernd Krauskopf, Hinke Osinga, Piotr Zgliczyński

# Plan of the presentation

- Toy example
- Cone conditions and covering relations
- Wall property and blender construction
- Computer assisted proof

---

- Greetings from Daniel Wilczak:


```
git clone https://github.com/CAPDGroup/CAPDexercises
```

# Toy example

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$$

diffeomorphism

- $x$  – strong expanding
- $y_u$  – weak expanding
- $y_s$  – contracting




# Toy example

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$$

diffeomorphism

- $x$  – strong expanding
- $y_u$  – weak expanding
- $y_s$  – contracting




# Toy example


$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$$

diffeomorphism

- $x$  – strong expanding
- $y_u$  – weak expanding
- $y_s$  – contracting



preimage of  $M$




$$\mathcal{G}(U) := f^{-1}(U) \cap M$$

# Toy example


$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$$

diffeomorphism

- $x$  – strong expanding
- $y_u$  – weak expanding
- $y_s$  – contracting



second preimage of  $M$



$$\mathcal{G}(U) := f^{-1}(U) \cap M$$

# Toy example


$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$$

diffeomorphism


- $x$  – strong expanding
- $y_u$  – weak expanding
- $y_s$  – contracting



third preimage of  $M$



second preimage of  $M$



$$\mathcal{G}(U) := f^{-1}(U) \cap M$$

# Toy example


$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$$

diffeomorphism


- $x$  – strong expanding
- $y_u$  – weak expanding
- $y_s$  – contracting



third preimage of  $M$



in the limit



$$\mathcal{G}(U) := f^{-1}(U) \cap M$$

$$\Lambda = \text{Inv}(f, M)$$

$$W^s(\Lambda) = \lim_{n \rightarrow \infty} \mathcal{G}^n(M)$$

$W^s(\Lambda)$  has one dimensional bundles, and yet behaves as a plane.



# Toy example


$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$$

diffeomorphism


- $x$  – strong expanding
- $y_u$  – weak expanding
- $y_s$  – contracting



third preimage of  $M$



in the limit



$$\mathcal{G}(U) := f^{-1}(U) \cap M$$

$$\Lambda = \text{Inv}(f, M)$$

$$W^s(\Lambda) = \lim_{n \rightarrow \infty} \mathcal{G}^n(M)$$

[BCDW] Bonatti, Crovisier, Díaz, Wilkinson, What is . . . a blender? Notices Amer. Math. Soc. (2016)

# Hyperbolic set

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

## Definition

We say that  $\Lambda$  is a hyperbolic set iff

$$T_z \Lambda = E_z^u \oplus E_z^s, \quad u + s = n$$


$$Df(z)E_z^u = E_{f(z)}^u$$

$$Df(z)E_z^s = E_{f(z)}^s$$

$$\|Df^n(z)v\| < c\lambda^n \|v\| \quad \text{for } v \in E_z^s$$

$$\|Df^{-n}(z)v\| < c\lambda^n \|v\| \quad \text{for } v \in E_z^u$$

where  $z \in \Lambda$  and  $\lambda \in (0, 1)$ .



$$\Lambda = \text{Inv}(f, M)$$

$$u = 1 \quad s = 1$$

# Hyperbolic set

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

## Definition

We say that  $\Lambda$  is a hyperbolic set iff

$$T_z \Lambda = E_z^u \oplus E_z^s, \quad u + s = n$$


$$Df(z)E_z^u = E_{f(z)}^u$$

$$Df(z)E_z^s = E_{f(z)}^s$$

$$\|Df^n(z)v\| < c\lambda^n \|v\| \quad \text{for } v \in E_z^s$$

$$\|Df^{-n}(z)v\| < c\lambda^n \|v\| \quad \text{for } v \in E_z^u$$

where  $z \in \Lambda$  and  $\lambda \in (0, 1)$ .



$$\Lambda = \text{Inv}(f, M)$$

$$u = 2 \quad s = 1$$

# Hyperbolic set

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

## Theorem


Let  $\Lambda = \text{Inv}(f, \bigcup N_i)$  and let

$$\mathcal{K} = \text{diag}(Id_u, -Id_s).$$

If for  $z \in f^{-1}(N_j) \cap N_i$

$$Df(z)^T \mathcal{K} Df(z) - \mathcal{K}$$

is positive definite, then  $\Lambda$  is a hyperbolic set.



$$\Lambda = \text{Inv}(f, M)$$

$$u = 2 \quad s = 1$$

[W] Wilczak, Uniformly hyperbolic attractor of the Smale-Williams type for a Poincaré map in the Kuznetsov system, SIADS (2010)

# Blender

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$


## Definition

We say that a hyperbolic set  $\Lambda$  is a  $k$ -blender if  $s < k$  and there exists an open set  $W$  of  $n - k$  dimensional surfaces such that

$$h \cap W_{\Lambda}^s \neq \emptyset \quad \forall h \in W.$$

Intuition:

- $W_{\Lambda}^s$  is 's-dimensional', but 'behaves as' a  $k$ -dimensional surface.



$$\Lambda = \text{Inv}(f, M)$$

$$u = 2 \quad s = 1$$

# Blender

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$


## Definition

We say that a hyperbolic set  $\Lambda$  is a  $k$ -blender if  $s < k$  and there exists an open set  $W$  of  $n - k$  dimensional surfaces such that

$$h \cap W_{\Lambda}^s \neq \emptyset \quad \forall h \in W.$$

Intuition:

- $W_{\Lambda}^s$  is 's-dimensional', but 'behaves as' a  $k$ -dimensional surface.



a 2-blender



$$\Lambda = \text{Inv}(f, M)$$

$$u = 2 \quad s = 1$$

# Covering relations


## Definition

We say that  $N_i \Rightarrow N_j$  iff



In the setting of the blender construction:

- 'Topological entry' coordinate  $y = (y_s, y_u)$  includes  $y_u$ , which is expanding.



# Cone conditions




$$C(z) = \{q : \|\pi_x(q - z)\| \geq \|\pi_y(q - z)\|\}$$

## Definition

$f$  satisfies cone conditions iff

$$f(C(z)) \subset C(f(z))$$

Notation:  $N \xrightarrow{cc} M$  means covering and cone conditions.










# horizontal discs

## Definition

Let  $N = B_x \times B_y$ . We say that  $h : B_x \rightarrow N$  is a horizontal disc in  $N$  iff

- $h(x) = (x, \pi_y h(x))$
- $h(B_x) \subset C(h(x))$






# horizontal discs

## Definition

Let  $N = B_x \times B_y$ . We say that  $h : B_x \rightarrow N$  is a horizontal disc in  $N$  iff

- $h(x) = (x, \pi_y h(x))$
- $h(B_x) \subset C(h(x))$






# horizontal discs

## Definition

Let  $N = B_x \times B_y$ . We say that  $h : B_x \rightarrow N$  is a horizontal disc in  $N$  iff

- $h(x) = (x, \pi_y h(x))$
- $h(B_x) \subset C(h(x))$






# horizontal discs

## Definition

Let  $N = B_x \times B_y$ . We say that  $h : B_x \rightarrow N$  is a horizontal disc in  $N$  iff

- $h(x) = (x, \pi_y h(x))$
- $h(B_x) \subset C(h(x))$



# Main result

$$M \in \{N_i\}_{i=1,\dots,m} \quad \Lambda = \text{Inv}(f, \bigcup N_i)$$


## Theorem

(A1) For every horizontal disc  $h \in M \exists i, j$   
 $h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$

(A3)  $\Lambda$  is hyperbolic


If (A1–A3) then  $\Lambda$  is a  $k$ -blender



$k$  is the dimension of the topological entry coordinate  $y = (y_u, y_s)$  and  $k > s$

below

$$\{N_i\} = \{N_1, N_2, M\}$$



# Main result

$$M \in \{N_i\}_{i=1,\dots,m} \quad \Lambda = \text{Inv}(f, \bigcup N_i)$$


## Theorem

(A1) For every horizontal disc  $h \in M \exists i, j$   
 $h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$

(A3)  $\Lambda$  is hyperbolic


If (A1–A3) then  $\Lambda$  is a  $k$ -blender



$k$  is the dimension of the topological entry coordinate  $y = (y_u, y_s)$  and  $k > s$

below

$$\{N_i\} = \{N_1, N_2, M\}$$



# Main result

$$M \in \{N_i\}_{i=1,\dots,m} \quad \Lambda = \text{Inv}(f, \bigcup N_i)$$


## Theorem

(A1) For every horizontal disc  $h \in M \exists i, j$   
 $h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$

(A3)  $\Lambda$  is hyperbolic


If (A1–A3) then  $\Lambda$  is a  $k$ -blender



$k$  is the dimension of the topological entry coordinate  $y = (y_u, y_s)$  and  $k > s$

below

$$\{N_i\} = \{N_1, N_2, M\}$$



# Main result

$$M \in \{N_i\}_{i=1,\dots,m} \quad \Lambda = \text{Inv}(f, \bigcup N_i)$$


## Theorem

(A1) For every horizontal disc  $h \in M \exists i, j$   
 $h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$

(A3)  $\Lambda$  is hyperbolic


If (A1–A3) then  $\Lambda$  is a  $k$ -blender



$k$  is the dimension of the topological entry coordinate  $y = (y_u, y_s)$  and  $k > s$

below

$$\{N_i\} = \{N_1, N_2, M\}$$





# Main result - proof

(A1) For every horizontal disc  $h \in M \exists i, j$   
 $h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$

$$\mathcal{G}(U) = f^{-1}(U) \cap \bigcup N_i$$


## Definition

$A \subset \bigcup N_i$  is a wall if for every  $h_i \in N_i$

$$h_i \cap A \neq \emptyset$$

## Lemma

If  $A$  is a wall and (A1–A2) then  $\mathcal{G}(A)$  is a wall.



# Main result - proof

(A1) For every horizontal disc  $h \in M \exists i, j$   
 $h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$

$$\mathcal{G}(U) = f^{-1}(U) \cap \bigcup N_i$$


## Definition

$A \subset \bigcup N_i$  is a wall if for every  $h_i \in N_i$

$$h_i \cap A \neq \emptyset$$

## Lemma

If  $A$  is a wall and  
 (A1–A2) then  $\mathcal{G}(A)$  is a  
 wall.



# Main result - proof

(A1) For every horizontal disc  $h \in M \exists i, j$

$$h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$   $\mathcal{G}(U) = f^{-1}(U) \cap \bigcup N_i$

## Definition


$A \subset \bigcup N_i$  is a wall if for every  $h_i \in N_i$

$$h_i \cap A \neq \emptyset$$

**Proof.**

## Lemma

If  $A$  is a wall and (A1–A2) then  $\mathcal{G}(A)$  is a wall.



[Z] Zgliczyński, Covering relations, cone conditions and the stable manifold theorem, JDE 2009

# Main result - proof

(A1) For every horizontal disc  $h \in M \exists i, j$

$$h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$   $\mathcal{G}(U) = f^{-1}(U) \cap \bigcup N_i$

## Definition


$A \subset \bigcup N_i$  is a wall if for every  $h_i \in N_i$

$$h_i \cap A \neq \emptyset$$

**Proof.**

## Lemma

If  $A$  is a wall and (A1–A2) then  $\mathcal{G}(A)$  is a wall.



[Z] Zgliczyński, Covering relations, cone conditions and the stable manifold theorem, JDE 2009

# Main result - proof

(A1) For every horizontal disc  $h \in M \exists i, j$

$$h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$   $\mathcal{G}(U) = f^{-1}(U) \cap \bigcup N_i$

## Definition


$A \subset \bigcup N_i$  is a wall if for every  $h_i \in N_i$

$$h_i \cap A \neq \emptyset$$

**Proof.**

## Lemma

If  $A$  is a wall and (A1–A2) then  $\mathcal{G}(A)$  is a wall.



[Z] Zgliczyński, Covering relations, cone conditions and the stable manifold theorem, JDE 2009

# Main result - proof

(A1) For every horizontal disc  $h \in M \exists i, j$

$$h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$   $\mathcal{G}(U) = f^{-1}(U) \cap \bigcup N_i$

## Definition


$A \subset \bigcup N_i$  is a wall if for every  $h_i \in N_i$

$$h_i \cap A \neq \emptyset$$

## Lemma

If  $A$  is a wall and (A1–A2) then  $\mathcal{G}(A)$  is a wall.

**Proof.**



[Z] Zgliczyński, Covering relations, cone conditions and the stable manifold theorem, JDE 2009

# Main result - proof

$$M \in \{N_i\}_{i=1,\dots,m} \quad \Lambda = \text{Inv}(f, \bigcup N_i)$$


## Theorem

(A1) For every horizontal disc  $h \in M \exists i, j$   
 $h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$

(A3)  $\Lambda$  is hyperbolic

If (A1–A3) then  $\Lambda$  is a  $k$ -blender



$k$  is the dimension of the topological entry coordinate  $y = (y_u, y_s)$  and  $k > s$

**Proof.**  $A_0 = \bigcup N_i$  is a wall.

$A_{j+1} = \mathcal{G}(A_j)$  are walls.

$A_j \rightarrow W_\Lambda^s$  as  $j \rightarrow \infty$  so  $W_\Lambda^s$  is a wall.

# Main result - proof

$$M \in \{N_i\}_{i=1,\dots,m} \quad \Lambda = \text{Inv}(f, \bigcup N_i)$$

## Theorem

(A1) For every horizontal disc  $h \in M \exists i, j$   
 $h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$


(A3)  $\Lambda$  is hyperbolic

If (A1–A3) then  $\Lambda$  is a  $k$ -blender

**Proof.**  $A_0 = \bigcup N_i$  is a wall.

$A_{j+1} = \mathcal{G}(A_j)$  are walls.

$A_j \rightarrow W_\Lambda^s$  as  $j \rightarrow \infty$  so  $W_\Lambda^s$  is a wall.



$k$  is the dimension of the topological entry coordinate  $y = (y_u, y_s)$  and  $k > s$



# Main result - proof

$$M \in \{N_i\}_{i=1,\dots,m} \quad \Lambda = \text{Inv}(f, \bigcup N_i)$$


## Theorem

(A1) For every horizontal disc  $h \in M \exists i, j$   
 $h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$

(A3)  $\Lambda$  is hyperbolic

If (A1–A3) then  $\Lambda$  is a  $k$ -blender



$k$  is the dimension of the topological entry coordinate  $y = (y_u, y_s)$  and  $k > s$

**Proof.**  $A_0 = \bigcup N_i$  is a wall.

$A_{j+1} = \mathcal{G}(A_j)$  are walls.

$A_j \rightarrow W_\Lambda^s$  as  $j \rightarrow \infty$  so  $W_\Lambda^s$  is a wall.



# Main result - some comments

$$M \in \{N_i\}_{i=1,\dots,m} \quad \Lambda = \text{Inv}(f, \bigcup N_i)$$

## Theorem


(A1) For every horizontal disc  $h \in M \exists i, j$

$$h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$

(A3)  $\Lambda$  is hyperbolic

If (A1–A3) then  $\Lambda$  is a  $k$ -blender




$k > s$  is the dimension of the topological entry coordinate  $y = (y_u, y_s)$

## Remarks:

- $\{N_i\}$  can consist of many sets
- We can generalise to  $\{M_l\}$
- Flexible framework

[BD] Bonatti, Díaz, Persistent nonhyperbolic transitive diffeomorphisms, Ann. of Math. (2), 143 (1996), 357–396,



# Main result - some comments

$$M \in \{N_i\}_{i=1,\dots,m} \quad \Lambda = \text{Inv}(f, \bigcup N_i)$$

## Theorem


(A1) For every horizontal disc  $h \in M \exists i, j$

$$h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$$

(A2)  $\forall i$  either  $N_i = M$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$

(A3)  $\Lambda$  is hyperbolic

If (A1–A3) then  $\Lambda$  is a  $k$ -blender





$k > s$  is the dimension of the topological entry coordinate  $y = (y_u, y_s)$

## Remarks:

- $\{N_i\}$  can consist of many sets
- We can generalise to  $\{M_l\}$
- Flexible framework

[BD] Bonatti, Díaz, Persistent nonhyperbolic transitive diffeomorphisms, Ann. of Math. (2), 143 (1996), 357–396,



# Main result - some comments

$$\{M_i\} \subset \{N_i\}_{i=1,\dots,m} \quad \Lambda = \text{Inv}(f, \bigcup N_i)$$

## Theorem


(A1) For every horizontal disc  $h \in M_i \exists i, j$

$$h_i = h \cap N_i \quad N_i \xrightarrow{cc} N_j$$

(A2)  $\forall i$  either  $N_i = M_i$  or  $\exists j$  s.t.  $N_i \xrightarrow{cc} N_j$

(A3)  $\Lambda$  is hyperbolic

If (A1–A3) then  $\Lambda$  is a  $k$ -blender




$k > s$  is the dimension of the topological entry coordinate  $y = (y_u, y_s)$

## Remarks:

- $\{N_i\}$  can consist of many sets
- We can generalise to  $\{M_i\}$
- Flexible framework

[BD] Bonatti, Díaz, Persistent nonhyperbolic transitive diffeomorphisms, Ann. of Math. (2), 143 (1996), 357–396,



# Example


Hénon family

$$f(x, y, z) = (y, \mu + y^2 + \beta x, \xi z + y)$$

$$\beta = 0.3, \quad \mu = -9.5.$$

## Theorem

*For every  $\xi \in [1.01, 1.125]$  the system has a 2-blender.*



# Example


Hénon family

$$f(x, y, z) = (y, \mu + y^2 + \beta x, \xi z + y)$$

$$\beta = 0.3, \quad \mu = -9.5.$$

## Theorem

For every  $\xi \in [1.01, 1.125]$  the system has a 2-blender.



$$N_{00} \xRightarrow{cc} N_{01} \xRightarrow{cc} N_{02} \xRightarrow{cc} N_{03} \xRightarrow{cc} M$$

# Example


Hénon family

$$f(x, y, z) = (y, \mu + y^2 + \beta x, \xi z + y)$$

$$\beta = 0.3, \quad \mu = -9.5.$$

## Theorem

For every  $\xi \in [1.01, 1.125]$  the system has a 2-blender.



$$N_{00} \xrightarrow{cc} N_{01} \xrightarrow{cc} N_{02} \xrightarrow{cc} N_{03} \xrightarrow{cc} M$$

# Example


Hénon family

$$f(x, y, z) = (y, \mu + y^2 + \beta x, \xi z + y)$$

$$\beta = 0.3, \quad \mu = -9.5.$$

## Theorem

*For every  $\xi \in [1.01, 1.125]$  the system has a 2-blender.*



$$N_{00} \xrightarrow{cc} N_{01} \xrightarrow{cc} N_{02} \xrightarrow{cc} N_{03} \xrightarrow{cc} M$$



# Example


Hénon family

$$f(x, y, z) = (y, \mu + y^2 + \beta x, \xi z + y)$$

$$\beta = 0.3, \quad \mu = -9.5.$$

## Theorem

*For every  $\xi \in [1.01, 1.125]$  the system has a 2-blender.*



$$N_{10} \xrightarrow{CC} N_{11} \xrightarrow{CC} N_{12} \xrightarrow{CC} N_{13} \xrightarrow{CC} N_{14} \xrightarrow{CC} M$$

# Example


Hénon family

$$f(x, y, z) = (y, \mu + y^2 + \beta x, \xi z + y)$$

$$\beta = 0.3, \quad \mu = -9.5.$$

## Theorem

*For every  $\xi \in [1.01, 1.125]$  the system has a 2-blender.*



$$N_{10} \xrightarrow{cc} N_{11} \xrightarrow{cc} N_{12} \xrightarrow{cc} N_{13} \xrightarrow{cc} N_{14} \xrightarrow{cc} M$$

# Example


Hénon family

$$f(x, y, z) = (y, \mu + y^2 + \beta x, \xi z + y)$$

$$\beta = 0.3, \quad \mu = -9.5.$$

## Theorem

*For every  $\xi \in [1.01, 1.125]$  the system has a 2-blender.*



$$N_{10} \xrightarrow{cc} N_{11} \xrightarrow{cc} N_{12} \xrightarrow{cc} N_{13} \xrightarrow{cc} N_{14} \xrightarrow{cc} M$$

# Example


Hénon family

$$f(x, y, z) = (y, \mu + y^2 + \beta x, \xi z + y)$$

$$\beta = 0.3, \quad \mu = -9.5.$$

## Theorem

For every  $\xi \in [1.01, 1.125]$  the system has a 2-blender.



$$N_{00} \xRightarrow{cc} N_{01} \xRightarrow{cc} N_{02} \xRightarrow{cc} N_{03} \xRightarrow{cc} M$$

# Example


Hénon family

$$f(x, y, z) = (y, \mu + y^2 + \beta x, \xi z + y)$$

$$\beta = 0.3, \quad \mu = -9.5.$$

## Theorem

*For every  $\xi \in [1.01, 1.125]$  the system has a 2-blender.*



$$N_{00} \xRightarrow{cc} N_{01} \xRightarrow{cc} N_{02} \xRightarrow{cc} N_{03} \xRightarrow{cc} M$$

# Example


Hénon family

$$f(x, y, z) = (y, \mu + y^2 + \beta x, \xi z + y)$$

$$\beta = 0.3, \quad \mu = -9.5.$$

## Theorem

For every  $\xi \in [1.01, 1.125]$  the system has a 2-blender.



$$N_{10} \xrightarrow{cc} N_{11} \xrightarrow{cc} N_{12} \xrightarrow{cc} N_{13} \xrightarrow{cc} N_{14} \xrightarrow{cc} M$$

# Example


Hénon family

$$f(x, y, z) = (y, \mu + y^2 + \beta x, \xi z + y)$$

$$\beta = 0.3, \quad \mu = -9.5.$$

## Theorem


For every  $\xi \in [1.01, 1.125]$  the system has a 2-blender.



$$N_{10} \xrightarrow{cc} N_{11} \xrightarrow{cc} N_{12} \xrightarrow{cc} N_{13} \xrightarrow{cc} N_{14} \xrightarrow{cc} M$$

## Closing remarks

- Blenders follow from covering relations and cone conditions
- Conditions (A1)–(A3) are verifiable using computer assisted proofs
- Flexible framework



### Friday: Heterodimensional cycles

[CKOZ] MC, B. Krauskopf, H. Osinga, P. Zgliczyński, Characterising blenders via covering relations and cone conditions. arXiv